

APPLICATION GUIDE FOR:

CALAMANSI

(*Citrofortunella microcarpa*)



Dilution:

- Mix 1 liter of SG7 with 100 liters of water. (1:100)
- Use unchlorinated water when mixing SG7. (Chlorine can kill microbes in the soil that may be beneficial to plants.)

Dosage:

- 1 ha = 2 liters of SG7 + 200 liters of water

Total SG7:

- 16 - 18 liters per hectare

IMPORTANT!

- Calamansi seedlings are ready for transplanting when they have grown to around 3 feet tall (approximately 1–12 months old).
- Transplant seedlings around June or July to ensure that water for irrigation is in adequate supply.
- Apply SG7 early in the morning, between 5–9 a.m. (maximum stomatal opening), or late in the afternoon, between 4–6 p.m. (extended stomatal opening).
- As soon as fruits start to develop from the flowers, stop application of SG7 to ensure that fruits are free of residual deposits.
- If cropping started from seedlings, the plant will reach its flowering stage in 18–24 months and its fruiting stage in another 10–12 months.

METHOD & FREQUENCY OF APPLICATION:

SEED, SEEDLING, AND SOIL TREATMENT:

For seeds - 1-2 hours before planting

For seedlings - 1-2 hours before transplanting

For soil - 1 - 3 days before planting or transplanting

- Dosage: 2L per hectare or 160 ml per 16L tank load
- Method (seeds): Spray mixture and let dry
- Method (seedlings): Spray or soak and let dry
- Method (soil): Apply directly to the soil until moist

1st APPLICATION:

10 - 15 days before planting or transplanting

- Dosage: 2L per hectare or 160 ml per 16L tank load
- Method: Apply directly to the soil (moist to dry)

SUCCESSING APPLICATION:

30 days after previous application with a maximum of 6–7 plant applications

- Dosage: 2L per hectare or 160 ml per 16L tank load
- Method: Apply directly to the plant

LAST APPLICATION:

10–15 days before flowering (approximately 9–10 months after transplanting)

- Dosage: 2L per hectare or 160 ml per 16L tank load
- Method: Apply directly to the plant